arXiv:1511.08075v1 [math.CA] 25 Nov 2015

ON WEAK*-CONVERGENCE IN THE LOCALIZED HARDY SPACES $H^1_\rho(\mathcal{X})$ AND ITS APPLICATION

DINH THANH DUC, HA DUY HUNG AND LUONG DANG KY *

Abstract. Let $(\mathcal{X}, d, \mu)$ be a complete RD-space. Let ρ be an admissible function on $\mathcal{X}$, which means that ρ is a positive function on $\mathcal{X}$ and there exist positive constants C_0 and k_0 such that, for any $x, y \in \mathcal{X}$,

$$\rho(y) \le C_0[\rho(x)]^{1/(1+k_0)}[\rho(x) + d(x,y)]^{k_0/(1+k_0)}.$$

In this paper, we define a space $VMO_\rho(\mathcal{X})$ and show that it is the predual of the localized Hardy space $H^1_\rho(\mathcal{X})$ introduced by Yang and Zhou [14]. Then we prove a version of the classical theorem of Jones and Journé [7] on weak*-convergence in $H^1_\rho(\mathcal{X})$. As an application, we give an atomic characterization of $H^1_\rho(\mathcal{X})$.

1. Introduction

It is a well-known and classical result (see [2]) that the space $BMO(\mathbb{R}^n)$ is the dual of the Hardy space $H^1(\mathbb{R}^n)$ one of the few examples of separable, nonreflexive Banach space which is a dual space. In fact, let $C_c(\mathbb{R}^n)$ be the space of all continuous functions with compact support and denote by $VMO(\mathbb{R}^n)$ the closure of $C_c(\mathbb{R}^n)$ in $BMO(\mathbb{R}^n)$, Coifman and Weiss showed in [2] that $H^1(\mathbb{R}^n)$ is the dual space of $VMO(\mathbb{R}^n)$, which gives to $H^1(\mathbb{R}^n)$ a richer structure than $L^1(\mathbb{R}^n)$. For example, the classical Riesz transforms $\nabla(-\Delta)^{-1/2}$ are not bounded on $L^1(\mathbb{R}^n)$, but are bounded on $H^1(\mathbb{R}^n)$. In addition, the weak*-convergence is true in $H^1(\mathbb{R}^n)$ (see [7]), which is useful in the application of Hardy spaces to compensated compactness (see [1]) and in the study of commutators of singular integral operators (see [8, 10]). Let $L = -\Delta + V$ be a Schrödinger operator on $\mathbb{R}^n$, $n \ge 3$, where V is a nonnegative function, $V \neq 0$, and belongs to the reverse Hölder class $RH_{n/2}(\mathbb{R}^n)$. The Hardy space associated with the Schrödinger operator L, $H^1_L(\mathbb{R}^n)$, is then defined as the set of functions $f \in L^1(\mathbb{R}^n)$ such that $\|f\|_{H^1_L} := \|\mathcal{M}_L f\|_{L^1} < \infty$, where $\mathcal{M}_L f(x) := \sup_{t>0} |e^{-tL} f(x)|$. Recently, Ky [9] established that the weak*-convergence is true in $H^1_L(\mathbb{R}^n)$, which is useful in studying the endpoint estimates for commutators of singular integral operators related to L (see [10]).

2010 *Mathematics Subject Classification.* 42B35.

Key words and phrases. H^1, BMO, VMO, spaces of homogeneous type.

The first author is supported by Vietnam National Foundation for Science and Technology Development (Grant No. 101.02-2014.32)

*Corresponding author.

Let $(\mathcal{X}, d, \mu)$ be an RD-space, which means that $(\mathcal{X}, d, \mu)$ is a space of homogeneous type in the sense of Coifman-Weiss with the additional property that a reverse doubling property holds in $\mathcal{X}$ (see Section 2). Typical examples for such RD-spaces include Euclidean spaces, Heisenberg groups, Lie groups of polynomial growth, or more generally, Carnot-Carathéodory spaces with doubling measures. We refer to the seminal paper of Han, Müller and Yang [4] for a systematic study of the theory of function spaces in harmonic analysis on RD-spaces. Recently, Yang and Zhou [14] introduced and studied the theory of localized Hardy spaces $H^1_\rho(\mathcal{X})$ related to the admissible functions ρ. There, they showed that this theory has a wide range of applications in studying the theory of Hardy spaces associated with Schrödinger operators or degenerate Schrödinger operators on $\mathbb{R}^n$, or associated with sub-Laplace Schrödinger operators on Heisenberg groups or connected and simply connected nilpotent Lie groups, see [14, Section 5] for details.

Given a complete RD-space $(\mathcal{X}, d, \mu)$ and an admissible function ρ, we denote by $BMO_\rho(\mathcal{X})$ the dual space of $H^1_\rho(\mathcal{X})$ (see Section 2) and $VMO_\rho(\mathcal{X})$ the closure in the BMO_ρ-norm of the space $C_c(\mathcal{X})$ of all continuous functions with compact support. The aim of the present paper is to show that $H^1_\rho(\mathcal{X})$ is a dual space and that the weak*-convergence is true in $H^1_\rho(\mathcal{X})$. Our main results can be read as follows:

Theorem 1.1. *The space $H^1_\rho(\mathcal{X})$ is the dual of the space $VMO_\rho(\mathcal{X})$.*

Theorem 1.2. *Suppose that $\{f_j\}_{j\geq 1}$ is a bounded sequence in $H^1_\rho(\mathcal{X})$, and that $f_j(x) \to f(x)$ for almost every $x \in \mathcal{X}$. Then, $f \in H^1_\rho(\mathcal{X})$ and $\{f_j\}_{j\geq 1}$ weak*-converges to f, that is, for every $\varphi \in VMO_\rho(\mathcal{X})$, we have*

$$\lim_{j\to\infty} \int_{\mathcal{X}} f_j(x)\varphi(x)d\mu(x) = \int_{\mathcal{X}} f(x)\varphi(x)d\mu(x).$$

It should be pointed out that when $\mathcal{X} \equiv \mathbb{R}^n$, $n \geq 3$, and $\rho(x) \equiv \sup\{r > 0 : \frac{1}{r^{n-2}} \int_{B(x,r)} V(y)dy \leq 1\}$, where V is in the reverse Hölder class $RH_{n/2}(\mathbb{R}^n)$, Theorem 1.2 is just the main theorem in the paper of Ky [9, Theorem 1.1].

Throughout the whole paper, C denotes a positive geometric constant which is independent of the main parameters, but may change from line to line.

2. Preliminaries

Let d be a quasi-metric on a set $\mathcal{X}$, that is, d is a nonnegative function on $\mathcal{X} \times \mathcal{X}$ satisfying

(a) $d(x, y) = d(y, x)$,
(b) $d(x, y) > 0$ if and only if $x \neq y$,
(c) there exists a constant $\kappa \geq 1$ such that for all $x, y, z \in \mathcal{X}$,

$$d(x, z) \leq \kappa(d(x, y) + d(y, z)).$$

A trip $(\mathcal{X}, d, \mu)$ is called a *space of homogeneous type* in the sense of Coifman-Weiss if μ is a regular Borel measure satisfying *doubling property*, i.e. there exists a constant $C > 1$ such that for all $x \in \mathcal{X}$ and $r > 0$,

$$\mu(B(x, 2r)) \le C\mu(B(x, r)).$$

Remark 2.1. *By* [2, Theorem (3.2)], *we see that if* $(\mathcal{X}, d, \mu)$ *is a complete space of homogeneous type, then the closure of* B *is a compact set for all ball* $B \subset \mathcal{X}$.

Recall (see [4]) that a space of homogeneous type $(\mathcal{X}, d, \mu)$ is called an *RD-space* if it satisfies *reverse doubling property*, i.e. there exists a constant $C > 1$ such that

$$\mu(B(x, 2r)) \ge C\mu(B(x, r))$$

for all $x \in \mathcal{X}$ and $r \in (0, \mathrm{diam}(\mathcal{X})/2)$, where $\mathrm{diam}(\mathcal{X}) := \sup_{x,y\in\mathcal{X}} d(x, y)$.

Here and what in follows, for $x, y \in \mathcal{X}$ and $r > 0$, we denote $V_r(x) := \mu(B(x, r))$ and $V(x, y) := \mu(B(x, d(x, y)))$.

Definition 2.1. *Let* $x_0 \in \mathcal{X}$, $r > 0$, $0 < \beta \le 1$ *and* $\gamma > 0$. *A function* f *is said to belong to the space of test functions,* $\mathcal{G}(x_0, r, \beta, \gamma)$, *if there exists a positive constant* C_f *such that*

(i) $|f(x)| \le C_f \frac{1}{V_r(x_0)+V(x_0,x)} \Big(\frac{r}{r+d(x_0,x)}\Big)^\gamma$ *for all* $x \in \mathcal{X}$;

(ii) $|f(x) - f(y)| \le C_f \Big(\frac{d(x,y)}{r+d(x_0,x)}\Big)^\beta \frac{1}{V_r(x_0)+V(x_0,x)} \Big(\frac{r}{r+d(x_0,x)}\Big)^\gamma$ *for all* $x, y \in \mathcal{X}$ *satisfying that* $d(x, y) \le \frac{r+d(x_0,x)}{2\kappa}$.

For any $f \in \mathcal{G}(x_0, r, \beta, \gamma)$, *we define*

$$\|f\|_{\mathcal{G}(x_0,r,\beta,\gamma)} := \inf\{C_f : \text{(i) } \textit{and} \text{ (ii) } \textit{hold}\}.$$

Let ρ be a positive function on $\mathcal{X}$. Following Yang and Zhou [14], the function ρ is said to *be admissible* if there exist positive constants C_0 and k_0 such that, for any $x, y \in \mathcal{X}$,

$$\rho(y) \le C_0[\rho(x)]^{1/(1+k_0)}[\rho(x) + d(x, y)]^{k_0/(1+k_0)}.$$

Throughout the whole paper, we always assume that $\mathcal{X}$ is a complete RD-space with $\mu(\mathcal{X}) = \infty$, and ρ is an admissible function on $\mathcal{X}$. Also we fix $x_0 \in \mathcal{X}$.

In Definition 2.1, it is easy to see that $\mathcal{G}(x_0, 1, \beta, \gamma)$ is a Banach space. For simplicity, we write $\mathcal{G}(\beta, \gamma)$ instead of $\mathcal{G}(x_0, 1, \beta, \gamma)$. Let $\epsilon \in (0, 1]$ and $\beta, \gamma \in (0, \epsilon]$, we define the space $\mathcal{G}^\epsilon_0(\beta, \gamma)$ to be the completion of $\mathcal{G}(\epsilon, \epsilon)$ in $\mathcal{G}(\beta, \gamma)$, and denote by $(\mathcal{G}^\epsilon_0(\beta, \gamma))'$ the space of all continuous linear functionals on $\mathcal{G}^\epsilon_0(\beta, \gamma)$. We say that f is a *distribution* if f belongs to $(\mathcal{G}^\epsilon_0(\beta, \gamma))'$.

Remark that, for any $x \in \mathcal{X}$ and $r > 0$, one has $\mathcal{G}(x, r, \beta, \gamma) = \mathcal{G}(x_0, 1, \beta, \gamma)$ with equivalent norms, but of course the constants are depending on x and r.

Let f be a distribution in $(\mathcal{G}^\epsilon_0(\beta, \gamma))'$. We define *the grand maximal functions* $\mathcal{M}(f)$ and $\mathcal{M}_\rho(f)$ as following

$$\mathcal{M}(f)(x) := \sup\{|\langle f, \varphi\rangle| : \varphi \in \mathcal{G}^\epsilon_0(\beta, \gamma), \|\varphi\|_{\mathcal{G}(x,r,\beta,\gamma)} \le 1 \text{ for some } r > 0\},$$

$$\mathcal{M}_\rho(f)(x) := \sup\{|\langle f, \varphi\rangle| : \varphi \in \mathcal{G}^\epsilon_0(\beta, \gamma), \|\varphi\|_{\mathcal{G}(x,r,\beta,\gamma)} \le 1 \text{ for some } r \in (0, \rho(x))\}.$$

Definition 2.2. *Let $\epsilon \in (0,1)$ and $\beta, \gamma \in (0, \epsilon)$.*

(i) *The Hardy space $H^1(\mathcal{X})$ is defined by*

$$H^1(\mathcal{X}) = \{f \in (\mathcal{G}_0^\epsilon(\beta,\gamma))' : \|f\|_{H^1} := \|\mathcal{M}(f)\|_{L^1} < \infty\}.$$

(ii) *The Hardy space $H^1_\rho(\mathcal{X})$ is defined by*

$$H^1_\rho(\mathcal{X}) = \{f \in (\mathcal{G}_0^\epsilon(\beta,\gamma))' : \|f\|_{H^1_\rho} := \|\mathcal{M}_\rho(f)\|_{L^1} < \infty\}.$$

Remark 2.2. *It was established in* [3] *that the space $H^1(\mathcal{X})$ coincides with the atomic Hardy $H^1_{\rm at}(\mathcal{X})$ of Coifman and Weiss* [2]*. Moreover, for all $f \in H^1(X)$,*

$$\|f\|_{L^1} \le C\|f\|_{H^1_\rho} \le C\|f\|_{H^1}.$$

Recall (see [2]) that a function $f \in L^1_{\rm loc}(\mathcal{X})$ is said to be in $BMO(\mathcal{X})$ if

$$\|f\|_{BMO} := \sup_B \frac{1}{\mu(B)} \int_B \Big| f(x) - \frac{1}{\mu(B)} \int_B f(y) d\mu(y) \Big| d\mu(x) < \infty,$$

where the supremum is taken all over balls $B \subset \mathcal{X}$. Denote by $VMO(\mathcal{X})$ the closure in BMO norm of $C_c(\mathcal{X})$. The following is well-known (see [2]).

Theorem 2.1. (i) *The space $BMO(\mathcal{X})$ is the dual space of $H^1(\mathcal{X})$.*
(ii) *The space $H^1(\mathcal{X})$ is the dual space of $VMO(\mathcal{X})$.*

Definition 2.3. *Let ρ be an admissible function and $\mathcal{D} := \{B(x,r) \subset \mathcal{X} : r \ge \rho(x)\}$. A function $f \in L^1_{\rm loc}(\mathcal{X})$ is said to be in $BMO_\rho(\mathcal{X})$ if*

$$\|f\|_{BMO_\rho} := \|f\|_{BMO} + \sup_{B \in \mathcal{D}} \frac{1}{\mu(B)} \int_B |f(x)| d\mu(x) < \infty.$$

It was established in [13] that

Theorem 2.2. *The space $BMO_\rho(\mathcal{X})$ is the dual space of $H^1_\rho(\mathcal{X})$.*

3. Proof of Theorems 1.1 and 1.2

We begin by recalling the following (see [14, Proposition 3.1]).

Lemma 3.1. *Let ρ be an admissible function. Then, there exists a function $K_\rho : \mathcal{X} \times \mathcal{X} \to \mathbb{R}$ and a positive constant C such that*

(i) *$K_\rho(x,y) \ge 0$ for all $x, y \in \mathcal{X}$, and $K_\rho(x,y) = 0$ if $d(x,y) > C\min\{\rho(x), \rho(y)\}$;*
(ii) *$K_\rho(x,y) \le C \frac{1}{\mu(B(x,\rho(x))) + \mu(B(y,\rho(y)))}$ for all $x, y \in \mathcal{X}$;*
(iii) *$K_\rho(x,y) = K_\rho(y,x)$ for all $x, y \in \mathcal{X}$;*
(iv) *$|K_\rho(x,y) - K_\rho(x,y')| \le C \frac{d(y,y')}{\rho(x)} \frac{1}{\mu(B(x,\rho(x))) + \mu(B(y,\rho(y)))}$ for all $x, y, z \in \mathcal{X}$ with $d(y,y') \le [\rho(x) + d(x,y)]/2$;*

(v) *for any* $x, x', y, y' \in \mathcal{X}$ *satisfying* $d(x,x') \le [\rho(y)+d(x,y)]/3$ *and* $d(y,y') \le [\rho(x)+d(x,y)]/3$, *we have*

$$\Big|[K_\rho(x,y)-K_\rho(x,y')] - [K_\rho(x',y)-K_\rho(x',y')]\Big| \le C\frac{d(x,x')}{\rho(y)}\frac{d(y,y')}{\rho(x)}\frac{1}{\mu(B(x,\rho(x)))+\mu(B(y,\rho(y)))};$$

(vi) $\int_{\mathcal{X}} K_\rho(x,y)d\mu(x) = 1$ *for all* $y \in \mathcal{X}$.

Given a function f in $L^1(\mathcal{X})$, following [14], we define

$$K_\rho(f)(x) = \int_{\mathcal{X}} K_\rho(x,y)f(y)d\mu(y)$$

for all $x \in \mathcal{X}$. It follows directly from Lemma 3.1 that

$$\int_{\mathcal{X}} K_\rho(f)(x)g(x)d\mu(x) = \int_{\mathcal{X}} K_\rho(g)(x)f(x)d\mu(x) \tag{3.1}$$

for all $f \in L^1(\mathcal{X})$ and $g \in L^\infty(\mathcal{X})$. Moreover, by Remark 2.1,

$$K_\rho(\phi) \in C_c(\mathcal{X}) \text{ for all } \phi \in C_c(\mathcal{X}), \tag{3.2}$$

and, for any $x \in \mathcal{X}$, the function $\mathbb{K}_\rho(x,\cdot) : \mathcal{X} \to \mathbb{R}$, defined by

$$\mathbb{K}_\rho(x,z) := \int_{\mathcal{X}} K_\rho(x,y)K_\rho(y,z)d\mu(y), \tag{3.3}$$

is in $C_c(\mathcal{X})$. Remark that $K_\rho(K_\rho(f))(x) = \int_{\mathcal{X}} \mathbb{K}_\rho(x,z)f(z)d\mu(z)$.

The following lemma is due to Yang and Zhou [14].

Lemma 3.2. *There exists a positive constant* C *such that*

(i) *for any* $f \in L^1(\mathcal{X})$,

$$\|K_\rho(f)\|_{H^1_\rho} \le C\|f\|_{L^1};$$

(ii) *for any* $g \in H^1_\rho(\mathcal{X})$,

$$\|g - K_\rho(g)\|_{H^1} \le C\|g\|_{H^1_\rho}.$$

As a consequence of Lemma 3.2 and (3.1), for any $\phi \in C_c(\mathcal{X})$,

$$\|\phi - K_\rho(K_\rho(\phi))\|_{BMO_\rho} \le C\|\phi\|_{BMO}. \tag{3.4}$$

Now we are ready to give the proofs of Theorems 1.1 and 1.2.

Proof of Theorem 1.1. Since $VMO_\rho(\mathcal{X})$ is a subspace of $BMO_\rho(\mathcal{X})$, which is the dual space of $H^1_\rho(\mathcal{X})$, every function f in $H^1_\rho(\mathcal{X})$ determines a bounded linear functional on $VMO_\rho(\mathcal{X})$ of norm bounded by $\|f\|_{H^1_\rho}$.

Conversely, given a bounded linear functional L on $VMO_\rho(\mathcal{X})$. Then,

$$|L(\phi)| \le \|L\|\|\phi\|_{VMO_\rho} \le \|L\|\|\phi\|_{L^\infty}$$

for all $\phi \in C_c(\mathcal{X})$. This implies (see [12]) that there exists a finite signed Radon measure ν on $\mathcal{X}$ such that, for any $\phi \in C_c(\mathcal{X})$,

$$L(\phi) = \int_{\mathcal{X}} \phi(x) d\nu(x),$$

moreover, the total variation of ν, $|\nu|(\mathcal{X})$, is bounded by $\|L\|$. Therefore,

$$\|K_\rho(K_\rho(\nu))\|_{H^1_\rho} \le C\|K_\rho(\nu)\|_{L^1} \le C|\nu|(\mathcal{X}) \le C\|L\| \tag{3.5}$$

by Lemma 3.2, where $K_\rho(\nu)(x) := \int_{\mathcal{X}} K_\rho(x,y)d\nu(y)$ for all $x \in \mathcal{X}$.

On the other hand, by (3.4) and (3.2), we have

$$\begin{aligned} |(L - K_\rho(K_\rho(L)))(\phi)| &= |L(\phi - K_\rho(K_\rho(\phi)))| \\ &\le \|L\|\|\phi - K_\rho(K_\rho(\phi))\|_{VMO_\rho} \\ &\le C\|L\|\|\phi\|_{BMO} \end{aligned}$$

for all $\phi \in C_c(\mathcal{X})$, where $K_\rho(K_\rho(L))(\phi) := \int_{\mathcal{X}} K_\rho(K_\rho(\nu))(x)\phi(x)d\mu(x)$. Consequently, by Theorem 2.1(ii), there exists a function h belongs $H^1(\mathcal{X})$ such that $\|h\|_{H^1} \le C\|L\|$ and

$$(L - K_\rho(K_\rho(L)))(\phi) = \int_{\mathcal{X}} h(x)\phi(x)d\mu(x)$$

for all $\phi \in C_c(\mathcal{X})$. This, together with (3.5), allows us to conclude that

$$L(\phi) = \int_{\mathbb{R}^d} f(x)\phi(x)d\mu(x)$$

for all $\phi \in C_c(\mathcal{X})$, where $f := h + K_\rho(K_\rho(\nu))$ is in $H^1_\rho(\mathcal{X})$ and satisfies that $\|f\|_{H^1_\rho} \le \|h\|_{H^1_\rho} + \|K_\rho(K_\rho(\nu))\|_{H^1_\rho} \le C\|L\|$. The proof of Theorem 1.1 is thus completed. □

Proof of Theorem 1.2. Let $\{f_{n_k}\}_{k=1}^\infty$ be an arbitrary subsequence of $\{f_n\}_{n=1}^\infty$. As $\{f_{n_k}\}_{k=1}^\infty$ is a bounded sequence in $H^1_\rho(\mathcal{X})$, by Theorem 1.1 and the Banach-Alaoglu theorem, there exists a subsequence $\{f_{n_{k_j}}\}_{j=1}^\infty$ of $\{f_{n_k}\}_{k=1}^\infty$ such that $\{f_{n_{k_j}}\}_{j=1}^\infty$ weak*-converges to g for some $g \in H^1_\rho(\mathcal{X})$. Therefore, by (3.3), for any $x \in \mathcal{X}$,

$$\begin{aligned} \lim_{j\to\infty} K_\rho(K_\rho(f_{n_{k_j}}))(x) &= \lim_{j\to\infty} \int_{\mathcal{X}} \mathbb{K}_\rho(x,z) f_{n_{k_j}}(z) d\mu(z) \\ &= \int_{\mathcal{X}} \mathbb{K}_\rho(x,z) g(z) d\mu(z) = K_\rho(K_\rho(g))(x). \end{aligned}$$

This implies that $\lim_{j\to\infty}[f_{n_{k_j}}(x) - K_\rho(K_\rho(f_{n_{k_j}}))(x)] = f(x) - K_\rho(K_\rho(g))(x)$ for almost every $x \in \mathcal{X}$. Hence, by Lemma 3.2 and [5, Theorem 1.1],

$$\|f - K_\rho(K_\rho(g))\|_{H^1} \le \sup_{j\ge1} \|f_{n_{k_j}} - K_\rho(K_\rho(f_{n_{k_j}})\|_{H^1} \le C \sup_{j\ge1} \|f_{n_{k_j}}\|_{H^1_\rho} < \infty,$$

moreover,

$$\lim_{j\to\infty}\int_{\mathcal{X}}[f_{n_{k_j}}(x)-K_\rho(K_\rho(f_{n_{k_j}}))(x)]\phi(x)d\mu(x)=\int_{\mathcal{X}}[f(x)-K_\rho(K_\rho(g))(x)]\phi(x)d\mu(x)$$

for all $\phi\in C_c(\mathcal{X})$. As a consequence, we obtain that

$$\begin{aligned}\|f\|_{H^1_\rho} &\le \|f-K_\rho(K_\rho(g))\|_{H^1_\rho}+\|K_\rho(K_\rho(g))\|_{H^1_\rho}\\ &\le C\|f-K_\rho(K_\rho(g))\|_{H^1}+C\|g\|_{H^1_\rho}\\ &\le C\sup_{j\ge1}\|f_{n_{k_j}}\|_{H^1_\rho}<\infty,\end{aligned}$$

moreover, by $\{f_{n_{k_j}}\}_{j=1}^\infty$ weak*-converges to g in $H^1_\rho(\mathcal{X})$, (3.1) and (3.2),

$$\begin{aligned}&\lim_{j\to\infty}\int_{\mathcal{X}}f_{n_{k_j}}(x)\phi(x)d\mu(x)\\ =\ &\lim_{j\to\infty}\int_{\mathcal{X}}[f_{n_{k_j}}(x)-K_\rho(K_\rho(f_{n_{k_j}}))(x)]\phi(x)d\mu(x)+\lim_{j\to\infty}\int_{\mathcal{X}}f_{n_{k_j}}(x)K_\rho(K_\rho(\phi))(x)d\mu(x)\\ =\ &\int_{\mathcal{X}}[f(x)-K_\rho(K_\rho(g))(x)]\phi(x)d\mu(x)+\int_{\mathcal{X}}g(x)K_\rho(K_\rho(\phi))(x)d\mu(x)\\ =\ &\int_{\mathcal{X}}f(x)\phi(x)d\mu(x).\end{aligned}$$

This, by $\{f_{n_k}\}_{k=1}^\infty$ be an arbitrary subsequence of $\{f_n\}_{n=1}^\infty$, allows us to complete the proof of Theorem 1.2. □

4. An application

The purpose of this section is to give an atomic characterization of $H^1_\rho(\mathcal{X})$ by using Theorems 1.1 and 1.2. First, we define the concept of atoms of log-type.

Definition 4.1. *Given $1<q\le\infty$. A measurable function a is called an (H^1_ρ,q)-atom of log-type related to the ball $B(x_0,r)$ if*

(i) *supp* $a\subset B(x_0,r)$,
(ii) $\|a\|_{L^q(\mathcal{X})}\le[\mu(B(x_0,r))]^{1/q-1}$,
(iii) $\left|\int_{\mathcal{X}}a(x)d\mu(x)\right|\le\frac{1}{\log\left(e+\frac{\rho(x_0)}{r}\right)}$.

The main result in this section can be read as follows:

Theorem 4.1. *Let $1<q\le\infty$. A function f is in $H^1_\rho(\mathcal{X})$ if and only if it can be written as $f=\sum_j\lambda_j a_j$, where a_j are (H^1_ρ,q)-atoms of log-type and $\sum_j|\lambda_j|<\infty$.*

Moreover, there exists a constant $C > 0$ such that, for any $f \in H^1_\rho(\mathcal{X})$,

$$\|f\|_{H^1_\rho} \le C \inf\left\{\sum_j |\lambda_j| : f = \sum_j \lambda_j a_j\right\} \le C\|f\|_{H^1_\rho}.$$

Before giving the proof of Theorem 4.1, let us recall the definition of H^1_ρ atoms introduced by Yang and Zhou [14].

Definition 4.2. *Given $1 < q \le \infty$. A measurable function a is called an (H^1_ρ, q)-atom related to the ball $B(x_0, r)$ if $r < \rho(x_0)$ and*

(i) *supp $a \subset B(x_0, r)$,*
(ii) *$\|a\|_{L^q(\mathcal{X})} \le [\mu(B(x_0, r))]^{1/q-1}$,*
(iii) *if $r < \rho(x_0)/4$, then $\int_{\mathcal{X}} a(x)d\mu(x) = 0$.*

Remark 4.1. *If a is an (H^1_ρ, q)-atom, then $\frac{1}{\log(e+4)}a$ is an (H^1_ρ, q)-atom of log-type.*

Proof of Theorem 4.1. By Remark 4.1 and [14, Theorems 3.2], it suffices to prove that there exists a constant $C > 0$ such that if f can be written as $f = \sum_j \lambda_j a_j$, where a_j are (H^1_ρ, q)-atoms of log-type related to the balls $B(x_j, r_j)$ and $\sum_j |\lambda_j| < \infty$, then $\|f\|_{H^1_\rho} \le C \sum_j |\lambda_j|$. Since Theorem 1.2, we only need to prove that

$$\|a_j\|_{H^1_\rho} \le C$$

for all j. This is reduced to showing that, for any $\phi \in C_c(\mathcal{X})$,

$$\left|\int_{\mathcal{X}} a_j(x)\phi(x)d\mu(x)\right| \le C\|\phi\|_{BMO_\rho} \tag{4.1}$$

by Theorem 1.1. To prove (4.1), let us consider the following two cases:

(a) The case: $r_j \ge \rho(x_j)$. Then, by the Hölder inequality and [14, Lemma 2.2],

$$\begin{aligned}\left|\int_{\mathcal{X}} a_j(x)\phi(x)d\mu(x)\right| &\le \|a_j\|_{L^q(B(x_j,r_j))}\|\phi\|_{L^{q'}(B(x_j,r_j))} \\ &\le [\mu(B(x_j, r_j))]^{1/q-1}C[\mu(B(x_j, r_j))]^{1/q'}\|\phi\|_{BMO_\rho} \\ &\le C\|\phi\|_{BMO_\rho}\end{aligned}$$

where and hereafter $1/q' + 1/q = 1$.

(b) The case: $r_j < \rho(x_j)$. Then, by the Hölder inequality, [14, Lemma 2.2] and [11, Lemma 2.1],

$$\begin{aligned}\left|\int_{\mathcal{X}} a_j(x)\phi(x)d\mu(x)\right| &\le \left|\int_{\mathcal{X}} a_j(\phi-\phi_{B(x_j,r_j)})d\mu\right| + |\phi_{B(x_j,r_j)}|\left|\int_{\mathcal{X}} a_j d\mu\right| \\ &\le \|a_j\|_{L^q(B(x_j,r_j))}\|\phi-\phi_{B(x_j,r_j)}\|_{L^{q'}(B(x_j,r_j))} + C\|\phi\|_{BMO_\rho} \\ &\le [\mu(B(x_j,r_j))]^{1/q-1}C[\mu(B(x_j,r_j))]^{1/q'}\|\phi\|_{BMO_\rho} + C\|\phi\|_{BMO_\rho} \\ &\le C\|\phi\|_{BMO_\rho},\end{aligned}$$

where $\phi_{B(x_j,r_j)} := \frac{1}{\mu(B(x_j,r_j))}\int_{B(x_j,r_j)}\phi d\mu$. This ends the proof of Theorem 4.1.

□

Acknowledgements. The paper was completed when the third author was visiting to Vietnam Institute for Advanced Study in Mathematics (VIASM). He would like to thank the VIASM for financial support and hospitality.

References

[1] R. R. Coifman, P.-L. Lions, Y. Meyer and S. Semmes, Compensated compactness and Hardy spaces. J. Math. Pures Appl. (9) 72 (1993), no. 3, 247–286.
[2] R. R. Coifman and G. Weiss, Extensions of Hardy spaces and their use in analysis. Bull. Amer. Math. Soc. 83 (1977), no. 4, 569-645.
[3] L. Grafakos, L. Liu and D. Yang, Maximal function characterizations of Hardy spaces on RD-spaces and their applications. Sci. China Ser. A 51 (2008), no. 12, 2253–2284.
[4] Y. Han, D. Müller and D. Yang, A theory of Besov and TriebelLizorkin spaces on metric measure spaces modeled on Carnot-Carathéodory spaces, Abstr. Appl. Anal. (2008), art. ID 893409, 250 pp.
[5] H. D. Hung and L. D. Ky, On weak*-convergence in the Hardy space H^1 over spaces of homogeneous type, arXiv:1510.01019.
[6] H. D. Hung, D. Q. Huy and L. D. Ky, A note on weak*-convergence in $h^1(\mathbb{R}^d)$, arXiv:1510.03597.
[7] P. W. Jones and J-L. Journé, On weak convergence in $H^1(\mathbf{R}^d)$. Proc. Amer. Math. Soc. 120 (1994), no. 1, 137-138.
[8] L. D. Ky, Bilinear decompositions and commutators of singular integral operators. Trans. Amer. Math. Soc. 365 (2013), no. 6, 2931–2958.
[9] L. D. Ky, On weak*-convergence in $H^1_L(\mathbb{R}^d)$. Potential Anal. 39 (2013), no. 4, 355–368.
[10] L. D. Ky, Endpoint estimates for commutators of singular integrals related to Schrödinger operators, Rev. Mat. Iberoam. (to appear) or arXiv:1203.6335.
[11] H. Lin and D. Yang, Pointwise multipliers for localized Morrey-Campanato spaces on RD-spaces. Acta Math. Sci. Ser. B Engl. Ed. 34 (2014), no. 6, 1677–1694.
[12] H. L. Royden, Real analysis. Third edition. Macmillan Publishing Company, New York, 1988.
[13] D. Yang, D. Yang and Y. Zhou, Localized Morrey-Campanato spaces on metric measure spaces and applications to Schrödinger operators, Nagoya Math. J. 198 (2010), 77-119.
[14] D. Yang and Y. Zhou, Localized Hardy spaces H^1 related to admissible functions on RD-spaces and applications to Schrödinger operators. Trans. Amer. Math. Soc. 363 (2011), no. 3, 1197-1239.

Dinh Thanh Duc

Department of Mathematics, University of Quy Nhon, 170 An Duong Vuong, Quy Nhon, Binh Dinh, Vietnam

E-mail: dinhthanhduc@qnu.edu.vn

Ha Duy Hung

High School for Gifted Students, Hanoi National University of Education, 136 Xuan Thuy, Hanoi, Vietnam

E-mail: hunghaduy@gmail.com

Luong Dang Ky

Department of Mathematics, University of Quy Nhon, 170 An Duong Vuong, Quy Nhon, Binh Dinh, Vietnam

E-mail: luongdangky@qnu.edu.vn